

# OxyStar™ 2E



## Oxyfluorfen Herbicide

OxyStar™ 2E is a familiar formulation of a long used selective herbicide that offers economical and effective control of many troublesome weeds in cotton and soybeans.



### Key Benefits

- OxyStar™ 2E is a low cost alternative to more commonly used post applied products in cotton and soybeans
- Offers excellent residual control of troublesome weeds when used as a fallow bed or burndown treatment
- Labeled crops can be planted 7 days after burndown application
- OxyStar™ 2E offers both pre-emergence and post-emergence control of labeled weeds
- OxyStar™ 2E can be applied using either ground or air equipment
- OxyStar™ 2E is activated with as little as ¼ inch of rain or irrigation water

### Active Ingredients

Oxyfluorfen	22.3%
Other Ingredients	77.7%
<b>Total</b>	<b>100%</b>

**Contains 2.0 pounds of oxyfluorfen per gal.**

**Contains petroleum distillates.**

**Always read and follow label directions.**

OxyStar™ 2E is not registered for use in every state. Please check with Albaugh Inc. or your local State Agency for product registration status. EPA Reg. No. 42750 – 136 AD012809

## Cotton and Soybean\* Fallow Bed Applications\*\*

Weeds Controlled		Rate: 1 to 2 Pints /Acre
Buttercup, Smallflower	Groundsel, Common	Purslane, Common
Cheeseweed (Malva)	Henbit	Redmaids
Evening Primrose, Cutleaf ***	Ladysthumb	Rocket, London
Fiddleneck, Coast	Minerslettuce	Shepherdspurse
Filaree, Broadleaf	Mustard Species	Sida, Prickly
Filaree, Redstem	Nettle, Burning	Sowthistle, Annual
Geranium, Carolina	Oxalis	Velvetleaf (Wild Cotton)
Groundcherry, Cutleaf	Pigweed, Redroot	

\* Not labeled for use on soybean fallow beds in California

\*\* Soybeans or cotton may be planted 7 days following application and the fallow bed should be worked thoroughly to a depth of 2 inches prior to planting.

\*\*\* Requires maximum rate and or multiple applications for effective control

## Soybeans

Weeds Controlled Preemergence		Rate: 1.5 to 3.0 Pints
Groundcherry, Cutleaf	Nightshade, Black	Sida, Prickly (Teaweed)
Jimsonweed	Pigweed, Redroot	Smartweed, Pennsylvania
Lambsquarters, Common	Poinsettia, Wild	Sowthistle, Common
Nightshade, American Black	Shepardspurse	Velvetleaf

  

Weeds Controlled Postemergence	( Post Directed Application)	
Cocklebur, Common	Mustard, Wild	Sesbania, Hemp
Croton, Tropic	Nightshade, American Black	Shepherdspurse
Groundcherry, Cutleaf	Nightshade, Black	Sicklepod
Groundcherry, Wright	Nightshade, Hairy	Sida, Prickly (Teaweed)
Jimsonweed	Pigweed, Redroot	Smartweed, Pennsylvania
Lambsquarters, Common	Purslane, Common	Velvetleaf
Morningglory, Annual (up to 6 leaf)		

## Cotton

Weeds Controlled Postemergence	(Post Directed Application)	Labeled Crop
Cocklebur, Common	Nightshade, American Black	Sesbania, Hemp
Croton, Tropic	Nightshade, Black	Sicklepod
Groundcherry, Cutleaf	Nightshade, Hairy	Sida, Prickly (Teaweed)
Groundcherry, Wright	Pigweed, Redroot	Smartweed, Pennsylvania
Jimsonweed	Poinsettia, Wild	Velvetleaf
Lambsquarters, Common	Purslane, Common	
Morningglory, Annual (up to 6 leaf)		

Refer to the label for the specific rate for your crop and farming practices